

Scalar interactions and the polarizations of $B \rightarrow \phi K^*$

Chuan-Hung Chen^{a*} and Chao-Qiang Geng^{b†}

^a*Department of Physics, National Cheng-Kung University, Tainan, 701 Taiwan*

^b*Department of Physics, National Tsing-Hua University, Hsin-Chu, 300 Taiwan*

(Dated: February 2, 2008)

Abstract

We try to understand the polarization puzzle in $B \rightarrow \phi K^*$ decays with a simple Higgs model associated with flavor changing neutral current at tree level. The new interactions can effectively reduce the longitudinal polarization $|A_0|^2$. In particular, we find that if the couplings of b -quark in different chiralities to Higgs are the same, the transverse polarization $|A_\perp|^2$ can receive the largest contribution and its value can be as large as 30%. On the other hand, with opposite sign in the couplings, the other transverse polarization $|A_\parallel|^2$ is enhanced.

* Email: phychen@mail.ncku.edu.tw

† Email: geng@phys.nthu.edu.tw

In terms of naive helicity analysis, it is well known that the transverse polarizations of vector bosons are associated with their masses. It was expected that the partitions of vector meson polarizations in B decays should have the same behavior. As a result, the ratio of various polarizations in two-body B meson decays can be estimated to be

$$|A_0|^2 : |A_\perp|^2 : |A_\parallel|^2 \sim 1 : \frac{m_V^2}{M_B^2} : \frac{m_V^2}{M_B^2}, \quad (1)$$

where A_0 and $A_\parallel$ belong to the mixtures of S and D-wave decay amplitudes while $A_\perp$ the P-wave one, which satisfy the identity

$$\sum_{\lambda=0,\parallel,\perp} |A_\lambda|^2 = 1. \quad (2)$$

According to Eq. (1), it is believed that in B decays with light vector mesons $A_{\perp(\parallel)}$ are much smaller than A_0 . The expectation is confirmed by BELLE [1] and BABAR [2] in $B \rightarrow \rho\rho$ decays, in which the longitudinal parts occupy over 95%. Furthermore, when the final states include heavy vector mesons, transverse polarizations can be relatively large. The conjecture is verified in $B \rightarrow J/\Psi K^*$ decays [3, 4], in which the longitudinal contribution is only about 60%.

However, the rule in Eq. (1) seems to be broken in $B \rightarrow \phi K^*$ decays. From the recent measurements of BELLE [5] and BABAR [4, 6], summarized in the Table I, it is quite clear that the longitudinal polarizations of $B \rightarrow K^* \phi$ are only around 50%. To solve the anomalous polarizations, the authors of Refs. [7, 8, 9, 10, 11] have proposed some solutions by introducing proper mechanisms such as large annihilation effect due to $(S - P) \otimes (S + P)$ interactions [7], the enhanced transversality from transverse gluon emitted by $b \rightarrow sg^{(*)}$ [8], final state interactions [9, 10] and new sets of form factors [11]. All above proposals are related to the uncertainties of low energy QCD. The possible new physics effects are also studied in the literature [12]. In this paper, firstly we reexamine the branching and polarization fractions of $B \rightarrow \phi K^*$ in the framework of perturbative QCD (PQCD) by fixing the hard scale for the involving Wilson coefficients within the SM [13]. And then, we introduce a new type of scalar interactions, which allows flavor changing neutral current (FCNC) at tree level. We will display that the new interactions could explain the branching ratios (BRs) and various polarizations in $B \rightarrow \phi K^*$.

It is known that the decay amplitude of $B \rightarrow V_1 V_2$ with the helicity can be generally

TABLE I: The polarization fractions and relative phases for $B \rightarrow \phi K^*$.

Model	Polarization	BELLE	BABAR
$K^{*+}\phi$	$ A_0 ^2$	$0.52 \pm 0.08 \pm 0.03$	$0.46 \pm 0.12 \pm 0.03$
	$ A_\perp ^2$	$0.19 \pm 0.08 \pm 0.02$	
	$\phi_\parallel(rad)$	$2.10 \pm 0.28 \pm 0.04$	
	$\phi_\perp(rad)$	$2.31 \pm 0.20 \pm 0.07$	
$K^{*0}\phi$	$ A_0 ^2$	$0.45 \pm 0.05 \pm 0.02$	$0.52 \pm 0.05 \pm 0.02$
	$ A_\perp ^2$	$0.30 \pm 0.06 \pm 0.02$	$0.22 \pm 0.05 \pm 0.02$
	$\phi_\parallel(rad)$	$2.39 \pm 0.24 \pm 0.04$	$2.34^{+0.23}_{-0.20} \pm 0.05$
	$\phi_\perp(rad)$	$2.51 \pm 0.23 \pm 0.04$	$2.47 \pm 0.25 \pm 0.05$

parametrized as [14]

$$\mathcal{M}^{(\lambda)} = \epsilon_{1\mu}^*(\lambda) \epsilon_{2\nu}^*(\lambda) [a g^{\mu\nu} + b P_2^\mu P_1^\nu + i c \epsilon^{\mu\nu\alpha\beta} P_{1\alpha} P_{2\beta}] . \quad (3)$$

Consequently, the helicity amplitudes are given by

$$\begin{aligned} H_{00} &= \frac{-1}{2m_1 m_2} [M_B^2 - m_1^2 - m_2^2] a + 2M_B^2 p^2 b , \\ H_{\pm\pm} &= a \mp M_B p c , \end{aligned}$$

where p is the magnitude of vector meson momenta. Note that we can define the polarization amplitudes to be

$$A_0 = \frac{H_{00}}{(\sum_h |H_h|^2)^{1/2}}, \quad A_{\parallel(\perp)} = \frac{1}{\sqrt{2}(\sum_h |H_h|^2)^{1/2}} (H_{++} \pm H_{--}), \quad (4)$$

so that Eq. (2) is satisfied. The relative phases between A_λ are described by $\phi_{\parallel(\perp)} = \text{Arg}(A_{\parallel(\perp)}/A_0)$. In order to merge the results calculated by PQCD [13], we rewrite Eq. (3) as

$$\mathcal{M} = M_B^2 \mathcal{M}_L + M_B^2 \mathcal{M}_N \epsilon_{1T}^* \cdot \epsilon_{2T}^* + i \mathcal{M}_T \varepsilon^{\alpha\beta\gamma\rho} \epsilon_{1T}^* \epsilon_{2T}^* P_{1\gamma} P_{2\rho}.$$

In terms of the well known effective Hamiltonian for the inclusive $b \rightarrow ss\bar{s}$ process [15], the various transition matrix elements in $B \rightarrow \phi K^*$ are written as

$$\mathcal{M}_H = V_{tb} V_{ts}^* [f_\phi E_H + E_H^N + f_B A_H + A_H^N], \quad (5)$$

where $H = L, N$ and T , $f_{\phi(B)}$ is the decay constant of ϕ (B), and $E_H^{(N)}$ and $A_H^{(N)}$ denote the factorization (nonfactorization) contributions of emission and annihilation topologies, respectively. We note that the Wilson coefficients of weak interactions have been included in $\{E\}$ and $\{A\}$ and their explicit expressions can be found in Ref. [13]. For simplicity, we will fix the scale, estimated by the energy of the exchanged hard gluon, to be around $t = \sqrt{\bar{\Lambda}M_B}$ where $\bar{\Lambda} \sim M_B - m_b$ with m_b being the b-quark mass. The various contributions associated with different scales are shown in Table II. Here, we have neglected the values

TABLE II: Values (in units of 10^{-3}) of transition matrix elements associated with different hard scales of t (GeV).

t	E_L	E_L^N	A_L	E_N	E_N^N
2.0	-13.90	$0.37 + i0.37$	$1.37 - 8.05$	-2.09	$(-1.28 + i0.04)10^{-1}$
1.8	-14.91	$0.40 + i0.39$	$1.49 - i8.69$	-2.24	$(-1.41 + i0.04)10^{-1}$
1.6	-16.13	$0.43 + i0.42$	$1.62 - i9.50$	-2.42	$(-1.57 + i0.03)10^{-1}$
t	A_N	E_T	E_T^N	A_T	
2.0	$-2.07 + i3.15$	-4.08	$(-2.65 - i0.25)10^{-1}$	$-3.90 + i6.46$	
1.8	$-2.23 + i3.40$	-4.38	$(-2.92 - i0.29)10^{-1}$	$-4.22 + i6.98$	
1.6	$-2.44 + i3.71$	-4.73	$(-3.26 - i0.33)10^{-1}$	$-4.62 + i7.62$	

of A^N since they are much smaller than the others. In our numerical estimations, we have used $f_\phi = 0.237$ GeV, $f_\phi^T = 0.22$ GeV, $f_{K^*} = 0.22$ GeV, $f_{K^*}^T = 0.17$ GeV, $f_B = 0.19$ GeV, $m_\phi = 1.02$ GeV, $m_{K^*} = 0.89$ GeV and $M_B = 5.28$ GeV. The wave functions of ϕ and K^* are referred to the results of light-cone sum rules (LCSRs) [16]. Using the values of Table II, the BR and polarizations in $B_d \rightarrow \phi K^{*0}$ can be easily obtained. To illustrate the effects of nonfactorization and annihilation, we fix $t = 1.6$ GeV and we find that $(BR, |A_0|^2, |A_\perp|^2) = (8.1 \times 10^{-6}, 0.93, 0.03)|_{E_H}$, $(10.93 \times 10^{-6}, 0.73, 0.12)|_{E_H, A_H}$ and $(9.42 \times 10^{-6}, 0.62, 0.17)|_{E_H, A_H, E_H^N}$ for contributions with $\{E_H\}$, $\{E_H, A_H\}$ and $\{E_H, A_H, E_H^N\}$, respectively. From the results, we find that the effects of annihilation and nonfactorization can enhance $A_\perp$, but they are still not enough to explain the central values of data in Table I. For completeness, we present the results with different hard scales in Table III. From Table III, we note that the polarizations are stable in different scales. It is difficult to further reduce the longitudinal polarization without introducing new mechanism. As a

TABLE III: Branching ratios (in units of 10^{-6}), polarizations and relative phases with different hard scales of t (GeV) for $B_d \rightarrow \phi K^{*0}$ in the SM.

t	BR	$ A_0 ^2$	$ A_{\parallel} ^2$	$ A_{\perp} ^2$	$\phi_{\parallel}(rad)$	$\phi_{\perp}(rad)$
2.0	6.93	0.628	0.206	0.166	2.16	2.15
1.8	8.02	0.625	0.207	0.167	2.15	2.14
1.6	9.46	0.622	0.209	0.169	2.15	2.13

comparision, we also calculate the decay of $B^+ \rightarrow \rho^+ K^{*0}$ in the SM and, explicitly, we find that $(BR, |A_0|^2, |A_{\perp}|^2)|_{B^+ \rightarrow \rho^+ K^{*0}} = (14.69 \times 10^{-6}, 0.72, 0.13)|_{E_H, A_H, E_H^N}$. Note that the current experimental data of BABAR and BELLE for $(BR, |A_0|^2)$ are $((17.0_{-3.9}^{+3.5}) \times 10^{-6}, 0.79 \pm 0.09)$ [17] and $((8.9 \pm 1.7 \pm 1.2) \times 10^{-6}, 0.43 \pm 0.11_{-0.02}^{+0.05})$ [18], respectively, which are not consistent with each other. Due to these inconclusive results in $B \rightarrow \rho K^*$, in this study we regard the polarization anomaly happens only in the decays of $B \rightarrow \phi K^*$.

We now try to find out if there exists some kind of new interactions which can induce large transverse polarizations in $B \rightarrow \phi K^*$, but not in the others, such as $B \rightarrow \rho\rho$ and $B \rightarrow \rho K^*$. Naturally, one could try the scalar interactions in which the couplings between the scalar and fermions are proportional to the fermion masses (m_f). In these models, the down-quark pair production is expected to be one order of magnitude smaller than that of the strange-quark pair. However, as known, the couplings in one-Higgs-doublet and type I two-Higgs-doublet models are suppressed by m_f/m_W . Although there is an enhancement factor $\tan\beta$ in the type II Higgs model, the effects of the $b \rightarrow s$ flavor change (FC) transition are one-loop suppressed. In order to get large transverse polarizations in $B \rightarrow \phi K^*$, we consider a new type of scalar interactions in which FCNC at tree level is allowed. Our another reason to try scalar interactions is that the new contributions on transverse polarizations should avoid the light meson mass dependence or the power suppression of m_{ϕ}/M_B , , unlike the SM in which $\langle \phi | \bar{s} \gamma_{\mu} s | 0 \rangle = m_{\phi} \epsilon_{\phi\mu}$ arises. For an illustration, we consider the hadronic matrix element $\langle \phi K^* | \bar{b} s \bar{s} s | B \rangle$. To get the factorizable parts, we need do the Fierz transformation. Explicitly, we have

$$\langle \phi K^* | \bar{b} s \bar{s} s | B \rangle \propto \frac{1}{4N_c} \langle \phi | \bar{s} \sigma_{\mu\nu} s | 0 \rangle \langle K^* | \bar{b} \sigma^{\mu\nu} s | B \rangle + \dots, \quad (6)$$

where $N_c = 3$ is the color factor, the factor $1/4$ is from the Fierz transformation, $\sigma_{\mu\nu} =$

$i[\gamma^\mu, \gamma^\nu]/2$ and $\{\dots\}$ denotes contributions from other operators such as the vectors and axial-vectors, which are suppressed by a factor of m_ϕ/m_B at the amplitude level. Since the nonlocal structure of ϕ is related to the term $\not{d}_\phi T \not{P} \Phi_\phi^T$ with Φ_ϕ^T being the twist-2 ϕ meson wave function, the factor $\langle \phi | \bar{s} \sigma_{\mu\nu} s | 0 \rangle \propto \epsilon_\phi^\mu P^\nu - \epsilon_\phi^\nu P^\mu$ which is clearly independent of m_ϕ . Hence, the hadronic suppression of scalar interactions can be only from color factor and Fierz coefficients. Next, we will demonstrate that scalar interactions have important influence on $B \rightarrow \phi K^*$. Before introducing a specific model, we start from a general interaction with a scalar boson S , given by

$$\mathcal{L}_{eff} = (C_{bs} \bar{b} P_R s + C_{sb} \bar{s} P_R b + C_{ss} \bar{s} P_R s) S + H.c. \quad (7)$$

where $P_{L(R)} = 1 \mp \gamma_5$. Since we are not dealing with the CP problem, the parameters C_{ij} are regarded as real numbers. From Eq. (7) the effective interaction for the process of $b \rightarrow ss\bar{s}$ is derived to be

$$\mathcal{L}_{eff} = \frac{C_{ss}}{m_S^2} \bar{b} (C_{bs} P_R + C_{sb} P_L) s \bar{s} s, \quad (8)$$

where m_S is the mass of the scalar.

The new contribution to $B \rightarrow \phi K^*$ due to the scalar interaction are shown in Fig. 1, where (a) and (b) stand for the factorizable and nonfactorizable effects, respectively. Since we have assumed that the couplings of the scalar interaction to fermions are proportional to m_f , the annihilation topologies can be neglected due to the suppression of m_u/v or m_d/v , comparing to emission topologies (Fig. 1) associated with m_s/v . Similar to the SM case,




FIG. 1: Diagrams for hadronic transition matrix elements due to the scalar interaction of $\bar{b}(1 - \gamma_5)s \bar{s} s$ with (a) factorizable and (b) nonfactorizable contributions.

the decay amplitudes for various helicities could be written as

$$\mathcal{M}_H^{NP} = \frac{C_{ss} C_{sb}}{2N_c m_S^2} [f_\phi F_H + N_H], \quad (9)$$

where F_H and N_H are the factorizable and nonfactorizable effects, respectively. Here, for simplicity, we have only presented the contributions of C_{bs} . The result of C_{sb} can be obtained by changing the sign in $H = L$ and N . C_{sb} and C_{bs} have the same contributions for $H = T$. The explicit expressions of F_H and N_H are shown in Appendix I and their values are given in Table II.

TABLE IV: Values (in units of 10^{-2}) of transition matrix elements for scalar interactions.

F_L	N_L	F_N	N_N	F_T	F_N
9.74	$-2.85 + i0.26$	-6.91	$0.44 + i0.31$	-20.27	$-0.038 + i0.29$

For a specific model, we concentrate on the generalized two-Higgs-doublet model (Model III) and the corresponding Yukawa Lagrangian for down-type quarks is described by [19]

$$\mathcal{L}_Y^{(III)} = \eta_{ij}^D \bar{Q}_{iL} \Phi_1 D_{jR} + \xi_{ij}^D \bar{Q}_{iL} \Phi_2 D_{jR} + H.c. , \quad (10)$$

where the indices $i(j)$ represent the possible quark flavors and ξ_{ij}^D denote the allowed FC effects. The vacuum expectation values (VEVs) of neutral Higgs fields are denoted by $\langle \Phi_{1(2)}^0 \rangle = v_{1(2)}$. For convenience, we can choose a proper basis such that only one scalar field possesses the VEV. Hence, the new scalar fields could be chosen to be $\phi_1^0 = \cos \beta \Phi_1^0 + \sin \beta \Phi_2^0 = (v + H^0 + i\chi^0)/\sqrt{2}$ and $\phi_2^0 = -\sin \beta \Phi_1^0 + \cos \beta \Phi_2^0 = (H^1 + iH^2)/\sqrt{2}$, where $v = \sqrt{v_1^2 + v_2^2}$ is the vacuum expectation value (VEV) of ϕ_1^0 , $\cos \beta (\sin \beta) = v_{1(2)}/v$, $H^{0(1)}$ and H^2 are CP-even and CP-odd Higgs bosons, and χ^0 is Goldstone boson, respectively. Since H^0 and H^1 are not physical eigenstates, the mass eigenstates could be parametrized by a mixing angle α as $h_{SM}^0 = H^0 \cos \alpha + H^1 \sin \alpha$ and $h^0 = -H^0 \sin \alpha + H^1 \cos \alpha$. When α goes to zero, h_{SM}^0 becomes the SM Higgs. It is known that to get naturally small FCNC at tree level, one can use the ansatz [19, 20]

$$\xi_{ij}^D = \lambda_{ij} \frac{\sqrt{m_i m_j}}{v}. \quad (11)$$

It has been analyzed phenomenologically that the coupling λ_{sb} for the transition of $b \rightarrow s$ may not be small and it could be as large as $O(10)$ [19]. Besides the coupling λ_{sb} , for $b \rightarrow ss\bar{s}$, we also need the information on λ_{ss} , which is flavor conserved. To understand the order of magnitude on λ_{ss} , we refer to the case of type II model, in which the corresponding coupling $\bar{s}sH^1$ is $m_s \cos \alpha / (v \cos \beta)$, i.e., λ_{ss} is order of $\cos \alpha / \cos \beta$. In the scenario of a large

$\tan\beta = v_2/v_1$, λ_{ss} could be order of $\tan\beta \sim m_t/m_b$. We note that the large enhancement of λ_{ss} is natural only for the type II model. In our considered type III model, we shall set $\lambda_{ss} = O(100)$, which implies that the Higgs coupling to the strange quark $\xi_{ss}^D = \lambda_{ss}m_s/v$ is $O(10^{-2})$. Since we only concern the non-leptonic decays, it is clear that the value of $\lambda_{\mu\mu} = O(1)$ for the muonic coupling given in Ref. [19] can be relaxed and there are no stringent limits for λ_{ij} .

To estimate the influence of scalar interactions, we set $\lambda_{ss} = 90$, $\lambda_{sb} = 5$ and $m_H = 150$ GeV and take $\zeta = \lambda_{bs}/\lambda_{sb}$ as a variable. By using the results of Tables II and IV, we present the BR and polarizations of $B_d \rightarrow \phi K^{*0}$ for different values of ζ in Table V. In Fig. 2, we show BR and $|A_{\parallel(\perp)}|^2$ as functions of (a) λ_{ss} with $\zeta = 0.2$ and $m_H = 150$ GeV, (b) ζ with $\lambda_{ss} = 90$ and $m_H = 150$ GeV and (c) m_H with $\zeta = 0.2$ and $\lambda_{ss} = 90$, respectively. From

TABLE V: Branching ratios (in units of 10^{-6}), polarizations and relative phases of $B_d \rightarrow \phi K^{*0}$ by combining the results of Tables II and IV with $t = 1.6$ GeV, $m_H = 150$ GeV, $\lambda_{ss} = 90$ and $\lambda_{sb} = 5$.

ζ	BR	$ A_0 ^2$	$ A_{\parallel} ^2$	$ A_{\perp} ^2$	$\phi_{\parallel}(rad)$	$\phi_{\perp}(rad)$
-1.0	11.28	0.56	0.30	0.14	2.16	2.18
-0.6	11.25	0.56	0.27	0.17	2.24	2.23
-0.2	11.28	0.55	0.24	0.21	2.22	2.26
0.0	11.32	0.54	0.23	0.23	2.21	2.27
0.2	11.37	0.53	0.22	0.25	2.20	2.28
0.6	11.53	0.52	0.19	0.29	2.17	2.30
1.0	11.76	0.50	0.17	0.33	2.15	2.31

the results of Table V and Fig. 2, we find that $|A_{\perp}|^2$ increases (decreases) if $\zeta > 0$ (< 0). In particular, when $\zeta > 0.6$, $|A_{\perp}|^2$ can be as large as 30%. We remark that the contributions of λ_{sb} and λ_{bs} to $\mathcal{M}_{L,N}$ are opposite in sign but to $\mathcal{M}_T$ the same sign. Therefore, if we take $\lambda_{sb} = \lambda_{bs}$, the scalar interactions can only contribute to $|A_{\perp}|^2$. On the other hand, if $\lambda_{sb} = -\lambda_{bs}$, only $|A_{\parallel}|^2$ gets affected as shown in Table V.

In summary, we have studied how scalar interactions effectively affect the polarizations in $B \rightarrow \phi K^*$ decays. We have illustrated that for the scalar interactions with FCNC couplings of $C_{sb} = C_{bs} \sim 1.6 \times 10^{-2}$ and FC one of $C_{ss} \sim 4.7 \times 10^{-2}$, the longitudinal




FIG. 2: Branching ratio and $|A_{||(\perp)}|^2$ in $B_d \rightarrow \phi K^{*0}$ as functions of (a) λ_{ss} with $\zeta = 0.2$ and $m_H = 150$ GeV, (b) ζ with $\lambda_{ss} = 90$ and $m_H = 150$ GeV and (c) m_H with $\zeta = 0.2$ and $\lambda_{ss} = 90$, respectively. The solid, dashed and dotted curves stand for $t = 2.0, 1.8$ and 1.6 GeV, respectively.

polarization $|A_0|^2$ can be 50% and the transverse polarization $|A_\perp|^2$ 30%. We have also found that the sign of $\zeta = \lambda_{sb}/\lambda_{bs} = C_{sb}/C_{bs}$ controls the relative magnitudes of $A_{||}$ and $A_\perp$.

Acknowledgments

This work is supported in part by the National Science Council of R.O.C. under Grant #s: NSC-93-2112-M-006-010 and NSC-93-2112-M-007-014.

I. APPENDIX: DECAY AMPLITUDES FOR NEW SCALAR INTERACTIONS

The transition matrix elements of factorizable and nonfactorizable effects for the effective interaction $\bar{b}(1 - \gamma_5)s \bar{s}s$ are given as follows: the factorizable amplitudes with various helicities are

$$\begin{aligned}
 F_L = & 2\pi C_F M_B^2 \int_0^1 dx_1 dx_3 \int_0^\infty db_1 db_3 \Phi_B(x_1) \{ [(1 + x_3 - r_\phi) \Phi_{K^*}(x_3) \\
 & + r_{K^*}(1 - 2x_3 + r_\phi x_3) (\Phi^t(x_3) + \Phi_{K^*}^s(x_3))] E_e(t_e^{(1)}) h_e(x_1, x_3, b_1, b_3) \\
 & + [2r_{K^*}(1 - r_\phi) \Phi_{K^*}^s(x_3)] E_e(t_e^{(2)}) h_e(x_3, x_1, b_3, b_1) \}, \quad (12)
 \end{aligned}$$

$$\begin{aligned}
F_N = & -2\pi C_F M_B^2 \int_0^1 dx_1 dx_3 \int_0^\infty b_1 db_1 b_3 db_3 \Phi_B(x_1) \{ [(1+x_3-r_\phi) \Phi_{K^*}^T(x_3) \\
& + r_{K^*}(1-2x_3) (\Phi^v(x_3) + \Phi_{K^*}^a(x_3)) - r_\phi r_{K^*} ((2+x_3) \Phi_{K^*}^v(x_3) - x_3 \Phi_{K^*}^a(x_3))] \\
& \times E_e(t_e^{(1)}) h_e(x_1, x_3, b_1, b_3) + [r_{K^*}(1-r_\phi) (\Phi_{K^*}^v(x_3) + \Phi_{K^*}^a(x_3))] \\
& \times E_e(t_e^{(2)}) h_e(x_3, x_1, b_3, b_1) \}, \tag{13}
\end{aligned}$$

$$\begin{aligned}
F_T = & -4\pi C_F M_B^2 \int_0^1 dx_1 dx_3 \int_0^\infty b_1 db_1 b_3 db_3 \Phi_B(x_1) \{ [(1+x_3+r_\phi) \Phi_{K^*}^T(x_3) \\
& + r_{K^*}(1-2x_3) (\Phi^v(x_3) + \Phi_{K^*}^a(x_3)) - r_\phi r_{K^*} ((2+x_3) \Phi_{K^*}^a(x_3) - x_3 \Phi_{K^*}^v(x_3))] \\
& \times E_e(t_e^{(1)}) h_e(x_1, x_3, b_1, b_3) + [r_{K^*}(1-r_\phi) (\Phi_{K^*}^v(x_3) + \Phi_{K^*}^a(x_3))] \\
& \times E_e(t_e^{(2)}) h_e(x_3, x_1, b_3, b_1) \}, \tag{14}
\end{aligned}$$

where $\{\Phi\}$ denote the distribution amplitudes of ϕ and K^* mesons. We consider the effects up to twist-3. The hard function h_e and E_e factor are

$$\begin{aligned}
h_e(x_1, x_3, b_1, b_3) = & K_0(\sqrt{x_1 x_3} M_B b_1) S_t(x_3) [\theta(b_1 - b_3) K_0(\sqrt{x_3} M_B b_1) I_0(\sqrt{x_3} M_B b_3) \\
& + \theta(b_3 - b_1) K_0(\sqrt{x_3} M_B b_3) I_0(\sqrt{x_3} M_B b_1)], \\
E_e(t) = & \alpha_s(t) S_B(t) S_{K^*}(t).
\end{aligned}$$

The Sudakov factors for K^* and B mesons and threshold resummation factor are given by

$$\begin{aligned}
S_B = & \exp \left[-s(x_1 P_B^+, b_1) - 2 \int_{1/b_1}^t \frac{d\mu}{\mu} \gamma(\alpha_s(\mu)) \right], \\
S_{K^*} = & \exp \left[-s(x_3 P_3^+, b_3) - s((1-x_3) P_2^+, b_3) - 2 \int_{1/b_2}^t \frac{d\mu}{\mu} \gamma(\alpha_s(\mu)) \right], \\
S_t(x) = & \frac{2^{1+2c} \Gamma(3/2 + c)}{\sqrt{\pi} \Gamma(1+c) [x(1-x)]^c},
\end{aligned}$$

where $\gamma = -\alpha_s/\pi$ which is the quark anomalous dimension, the variables (b_1, b_2, b_3) are conjugate to the parton transverse momenta (k_{1T}, k_{2T}, k_{3T}) , $c = 0.4$ for $B \rightarrow \phi K^*$ decays, and the explicit expression for $s(x, b)$ can be found in Ref. [21]. The scale $t_e^{(1)}$ and $t_e^{(2)}$ are chosen by

$$\begin{aligned}
t_e^{(1)} = & \max(\sqrt{x_3} M_B, 1/b_1, 1/b_3), \\
t_e^{(2)} = & \max(\sqrt{x_1} M_B, 1/b_1, 1/b_3).
\end{aligned}$$

The nonfactorizable amplitudes with various helicities are given as

$$\begin{aligned}
N_L = & -4\pi C_F M_B^2 \sqrt{2N_c} \int_0^1 d[x] \int_0^\infty b_1 db_1 b_2 db_2 \Phi_B(x_1) \{ [x_2 \Phi_\phi(x_2) \Phi_{K^*}(x_3) \\
& + r_{K^*} x_3 (\Phi^t(x_3) - \Phi_{K^*}^s(x_3)) - r_\phi r_{K^*} \Phi_\phi^t(x_2) (0.5(x_2 + 3x_3) \Phi_{K^*}^t(x_3) + x_2 \Phi_{K^*}^s(x_3)) \\
& + 2r_\phi x_2 \Phi_\phi^t(x_2) \Phi_{K^*}(x_3) - r_\phi r_{K^*} ((x_2 - x_3) \Phi_{K^*}^t(x_3) + (x_2 + x_3) \Phi_{K^*}^s(x_3))] \\
& \times E_d(t_d^{(1)}) h_d^{(1)}(x_1, x_2, x_3, b_1, b_2) - [2(1 - x_2 + x_3) \Phi_\phi(x_2) \Phi_{K^*}(x_3) \\
& + r_{K^*} x_3 \Phi_\phi(x_2) (\Phi^t(x_3) + \Phi_{K^*}^s(x_3)) - r_\phi (1 - x_2) (\Phi_\phi^t(x_2) - \Phi_\phi^s(x_2)) \Phi_{K^*}(x_3) \\
& + 2r_\phi r_{K^*} (1 - x_2 + x_3) (\Phi_\phi^t(x_2) \Phi_{K^*}^t(x_3) - \Phi_\phi^s(x_2) \Phi_{K^*}^s(x_3))] \\
& \times E_d(t_d^{(2)}) h_d^{(1)}(x_1, x_2, x_3, b_1, b_2) \}, \tag{15}
\end{aligned}$$

$$\begin{aligned}
N_N = & -4\pi C_F M_B^2 \sqrt{2N_c} \int_0^1 d[x] \int_0^\infty b_1 db_1 b_2 db_2 \Phi_B(x_1) \{ [-x_2 \Phi_\phi^T(x_2) \Phi_{K^*}^T(x_3) \\
& + r_\phi x_2 (\Phi_\phi^v(x_2) - \Phi_\phi^a(x_2)) \Phi_{K^*}^T(x_3)] E_d(t_d^{(1)}) h_d^{(1)}(x_1, x_2, x_3, b_1, b_2) \\
& + [(1 - x_2 + x_3) \Phi_\phi^T(x_2) \Phi_{K^*}^T(x_3) - r_{K^*} x_3 \Phi_\phi^T(x_2) (\Phi_{K^*}^v(x_3) + \Phi_{K^*}^a(x_3)) \\
& + r_\phi (1 - x_2) (\Phi_\phi^v(x_2) - \Phi_\phi^a(x_2)) \Phi_{K^*}^T(x_3) - 2r_\phi r_{K^*} (1 - x_2 + x_3) \\
& \times (\Phi_\phi^v(x_2) \Phi_{K^*}^v(x_3) - \Phi_\phi^a(x_2) \Phi_{K^*}^a(x_3))] E_d(t_d^{(2)}) h_d^{(2)}(x_1, x_2, x_3, b_1, b_2), \tag{16}
\end{aligned}$$

$$\begin{aligned}
N_T = & 8\pi C_F M_B^2 \sqrt{2N_c} \int_0^1 d[x] \int_0^\infty b_1 db_1 b_2 db_2 \Phi_B(x_1) \{ [x_2 \Phi_\phi^T(x_2) \Phi_{K^*}^T(x_3) \\
& + r_\phi x_2 (\Phi_\phi^v(x_2) - \Phi_\phi^a(x_2)) \Phi_{K^*}^T(x_3)] E_d(t_d^{(1)}) h_d^{(1)}(x_1, x_2, x_3, b_1, b_2) \\
& - [(1 - x_2 + x_3) \Phi_\phi^T(x_2) \Phi_{K^*}^T(x_3) - r_{K^*} x_3 \Phi_\phi^T(x_2) (\Phi_{K^*}^v(x_3) + \Phi_{K^*}^a(x_3)) \\
& - r_\phi (1 - x_2) (\Phi_\phi^v(x_2) - \Phi_\phi^a(x_2)) \Phi_{K^*}^T(x_3) + 2r_\phi r_{K^*} (1 - x_2 + x_3) \\
& \times (\Phi_\phi^v(x_2) \Phi_{K^*}^a(x_3) - \Phi_\phi^a(x_2) \Phi_{K^*}^v(x_3))] E_d(t_d^{(2)}) h_d^{(2)}(x_1, x_2, x_3, b_1, b_2), \tag{17}
\end{aligned}$$

where the Sudakov factor for the ϕ meson is given as

$$S_\phi = \exp \left[-s(x_2 P_2^+, b_2) - s((1 - x_2) P_2^+, b_2) - 2 \int_{1/b_2}^t \frac{d\mu}{\mu} \gamma(\alpha_s(\mu)) \right],$$

and the hard functions $h_d^{(j)}$ are

$$\begin{aligned}
h_d^{(j)} = & [\theta(b_1 - b_2) K_0(DM_B b_1) I_0(DM_B b_2) + \theta(b_2 - b_1) K_0(DM_B b_2) I_0(DM_B b_1)] \\
& \times \begin{cases} K_0(D_j M_B b_2) & \text{for } D_j^2 \geq 0, \\ \frac{i\pi}{2} H_0^{(1)}(\sqrt{|D_j^2|} M_B b_2) & \text{for } D_j^2 \leq 0 \end{cases}
\end{aligned}$$

with $D^2 = x_1 x_3$, $D_1^2 = (x_1 - x_2)x_3$ and $D_2^2 = -(1 - x_1 - x_2)x_3$. The scales $t_d^{(j)}$ are chosen by

$$t_d^{(1)} = \max(DM_B, \sqrt{|D_1^2|}M_B, 1/b_1, 1/b_2),$$

$$t_d^{(2)} = \max(DM_B, \sqrt{|D_2^2|}M_B, 1/b_1, 1/b_2).$$

-
- [1] BELLE Collaboration, J. Zhang *et al.*, Phys. Rev. Lett. **91**, 221801 (2003).
 - [2] BABAR Collaboration, B. Aubert *et al.*, Phys. Rev. Lett. **91**, 171802 (2003).
 - [3] BELLE Collaboration, K. Abe *et al.*, Phys. Lett. **B538**, 11 (2002); hep-ex/0408104.
 - [4] BABAR Collaboration, B. Aubert *et al.*, Phys. Rev. Lett. **87**, 241801 (2001).
 - [5] BELLE Collaboration, K. F. Chen, *et al.*, hep-ex/0503013.
 - [6] BABAR Collaboration, B. Aubert *et al.*, Phys. Rev. Lett. **93**, 231804 (2004).
 - [7] A. Kagan, Phys. Lett. **B601**, 151 (2004).
 - [8] W.S. Hou and M. Nagashima, hep-ph/0408007.
 - [9] P. Colangelo, F. De Fazio and T.N. Pham, Phys. Lett. **B597**, 291 (2004); M. Ladisa *et al.*, Phys. Rev. **D70**, 114025 (2004).
 - [10] H.Y. Cheng, C.K. Chua and A. Soni, Phys. Rev. **D71**, 014030 (2005).
 - [11] H.N. Li, hep-ph/0411305.
 - [12] A. Kagan, hep-ph/0407076; E. Alvarez *et al.*, Phys. Rev. **D70**, 115014 (2004); Y.D. Yang, R.M. Wang and G.R. Lu, hep-ph/0411211; A.K. Giri and R. Mohanta, hep-ph/0412107; P.K. Das and K.C. Yang, Phys. Rev. **D71**, 094002 (2005); C.S. Kim and Y.D. Yang, hep-ph/0412364.
 - [13] C.H. Chen, Y.Y. Keum and H.N. Li, Phys. Rev. **D66**, 054013 (2002).
 - [14] G. Valencia, Phys. Rev. **D39**, 3339 (1989); G. Kramer and W.F. Paimer, Phys. Rev. **D45**, 193 (1992).
 - [15] G. Buchalla, A.J. Buras and M.E. Lautenbacher, Rev. Mod. Phys. **68**, 1125 (1996).
 - [16] P. Ball *et al.*, Nucl. Phys. **B529**, 323 (1998).
 - [17] BABAR Collaboration, B. Aubert *et al.*, hep-ex/0408093.
 - [18] BELLE Collaboration, J. Zhang *et al.*, hep-ex/0505039.
 - [19] D. Atwood, L. Reina and A. Soni, Phys. Rev. **D55**, 3156 (1997).
 - [20] T.P. Cheng and M. Sher, Phys. Rev. **D35**, 3484 (1987); *ibid* **44**, 1461 (1991).

- [21] J. Botts and G. Sterman, Nucl. Phys. **B325**, 62 (1989); H.N. Li and G. Sterman, Nucl. Phys. **B381**, 129 (1992).